


ARPES on iron-based superconductors: leading role of $3d_{xz,yz}$ orbitals

D. Evtushinsky,

Institute for Solid State Research, IFW Dresden, Germany

I will start with a short overview of the available ARPES results on iron-based superconductors. Despite of the large variety of the observed Fermi surfaces [1-6], phenomenologically one can point out several common tendencies for iron high temperature superconductors — namely, band renormalization of factor of three at the Fermi level [3,6-8], often a two-gap behavior [2,7-10], noticeable electronic coupling to low-energy bosonic modes. Our recent results indicate clear correlation between the superconducting gap magnitude and orbital origin of the electronic states, with the largest gap solely for iron $3d_{xz,yz}$ orbitals.

Interestingly, the electron coupling to bosonic modes was found to be stronger also for $3d_{xz,yz}$ bands. Further analysis of the electronic structures of differently doped iron arsenides consistently points to the significance of $3d_{xz,yz}$ bands for superconductivity in these materials.



- [1] V. B. Zabolotnyy et al., *Nature* 457, 569 (2009).
- [2] D. V. Evtushinsky et al., *New J. Phys.* 11, 055069 (2009).
- [3] S. V. Borisenko et al., *Phys. Rev. Lett.* 105, 067002 (2010).
- [4] D. V. Evtushinsky et al., *J. Phys. Soc. Jpn.* 80 023710 (2011).
- [5] S. Thirupathaiah et al., *Phys. Rev. B* 86, 214508 (2012).
- [6] S. V. Borisenko et al., arXiv:1204.1316.
- [7] H. Ding et al., *Europhys. Lett.* 83, 47001 (2008).
- [8] D. V. Evtushinsky et al., arXiv:1204.2432.
- [9] D. V. Evtushinsky et al., *Phys. Rev. B* 79, 054517 (2009).
- [10] D. V. Evtushinsky et al., *Phys. Rev. B* 87, 094501 (2013).