

Electron-hole fluctuations driving the charge density wave phase transition in TiSe_2

Claude Monney

Fritz-Haber-Institut der Max Planck Gesellschaft, Berlin, Germany

Abstract:

There is a longstanding open question about the existence of a condensate of electron-hole pairs (excitons) in a way as it is well known for pairs of electrons, Cooper-pairs, in superconductivity. TiSe_2 exhibits an unusual temperature-dependence of transport experiments and a specific band configuration that has been related to the possible formation of excitons. Indeed, at the temperature of 200K, the layered, quasi-two-dimensional TiSe_2 undergoes a phase transition towards a charge density wave (CDW) phase, which is accompanied by a periodic lattice distortion (PLD) [1].

I propose the exciton condensate phase model [2] as the electronic origin of the CDW phase of TiSe_2 . This scenario is supported by the good agreement between measured angle-resolved photoemission spectroscopy data and photoemission intensity maps calculated by the spectral function within this model [3]. With the help of perturbation theory calculations, I will motivate that above 200K, strong electron-hole fluctuations are provoking an electronic instability towards the CDW phase [4]. Eventually, this instability drives the PLD via the electron-phonon coupling [5].

References:

- [1] F. J. Di Salvo *et al.*, Phys. Rev. B 14, 4321 (1976).
- [2] D. Jérôme *et al.*, Phys. Rev., 158 462 (1967).
- [3] H. Cercellier *et al.*, Phys. Rev. Lett. 99, 146403 (2007).
- [4] C. Monney *et al.*, New J. Phys. 14, 075026 (2012).
- [5] C. Monney *et al.*, Phys. Rev. Lett. 106, 106404 (2011).